


Cite this article: Rachmawati, M., Septianasari, V., Maulida, S., Apriadi, R., Suprpto H., Murdianto, W., & Saragih, B. (2026). Optimization and characterization of Tiwai (*Eleutherine americana* L. Merr) extract encapsulation for potential bioavailability of flavonoids. *Journal of Current Science and Technology*, 16(2), Article 172. <https://doi.org/10.59796/jcst.V16N2.2026.172>



## Optimization and Characterization of Tiwai (*Eleutherine americana* L. Merr.) Extract Encapsulation for Potential Bioavailability of Flavonoids

Maulida Rachmawati<sup>1</sup>, Vivie Septianasari<sup>2</sup>, Sarah Maulida<sup>2</sup>, Rimbawan Apriadi<sup>2</sup>, Hadi Suprpto<sup>2</sup>,  
Wiwit Murdianto<sup>2</sup>, and Bernatal Saragih<sup>1,2,\*</sup>

<sup>1</sup>Doctoral Program of Agricultural Science, Faculty of Agriculture, Mulawarman University,  
Kalimantan Timur 75117, Indonesia.

<sup>2</sup>Department of Agricultural Product Technology, Faculty of Agriculture, Mulawarman University,  
Kalimantan Timur 75117, Indonesia

\*Corresponding author; E-mail: [bernatalsaragih@faperta.unmul.ac.id](mailto:bernatalsaragih@faperta.unmul.ac.id)

Received 30 July 2025; Revised 10 October 2025; Accepted 28 October 2025; Published online 30 March 2026

### Abstract

Tiwai (*Eleutherine americana* L. Merr.) is a traditional medicinal plant from Kalimantan, rich in bioactive compounds, particularly flavonoids, which exhibit strong antioxidant activity and therapeutic benefits. However, flavonoids in free form tend to be unstable, making encapsulation necessary to enhance their stability, bioavailability, and shelf life. This study optimized the encapsulation of Tiwai extract using the spray drying method, with parameters such as the ratio of Tiwai extract to maltodextrin (ET:MD), chitosan concentration, and drying inlet temperature adjusted using Response Surface Methodology (RSM) with Box-Behnken Design (BBD). The study provided detailed results, including an encapsulation efficiency (EE) of 93.24%, loading capacity (LC) of 95.76%, total flavonoid content (TFC) of 69.23 mg QE/g, and antioxidant activity (DPPH) of 60.01 µg/mL. A significant discrepancy was found between the quercetin content determined by HPLC (148.30 µg/g) and the TFC obtained using the AlCl<sub>3</sub> method (69.23 mg QE/g). This difference is attributed to the different principles of the two methods, as the AlCl<sub>3</sub> method reacts with flavonoids and other antioxidant compounds, whereas HPLC specifically identifies quercetin. FTIR analysis confirmed successful encapsulation, showing changes in the functional groups of the encapsulated product. These findings suggest that the optimized Tiwai encapsulation formula has potential applications in functional foods and pharmaceuticals.

**Keywords:** antioxidant; encapsulation; flavonoids; spray drying; FTIR; HPLC

### 1. Introduction

Tiwai (*Eleutherine americana* L. Merr.), also known as Dayak onion, is a traditional medicinal plant widely found in Borneo, Indonesia. It is rich in bioactive compounds, especially flavonoids such as quercetin, kaempferol, and luteolin, which exhibit strong antioxidant, anti-inflammatory, and antibacterial properties. These flavonoids play a key role in the plant's bioactivity and offer various therapeutic benefits, such as treating hypertension, diabetes, and

skin infections (Adesina et al., 2016; Tiwari & Shukla, 2020; Kareem et al., 2022; Rachmawati et al., 2024). However, flavonoids in their free form are unstable, which limits their bioavailability and shelf life when used in functional foods or pharmaceuticals (Cocuron et al., 2019; Shahidi & Yeo, 2016). Therefore, encapsulation is required to improve their stability, bioavailability, and extend their shelf life in product formulations.

Encapsulation is a well-established method for protecting bioactive compounds such as flavonoids. Among the various techniques, spray drying is one of the most widely used methods. It works by converting liquid extracts into dry powder particles through rapid evaporation using a hot air stream. This process not only enhances the stability of the active compounds but also facilitates their incorporation into functional foods, nutraceuticals, and pharmaceutical formulations. Factors influencing the spray drying encapsulation process include the concentration of active ingredients, the type and concentration of coating materials, and the inlet temperature. These factors determine the efficiency, particle size, stability, and antioxidant activity of the final encapsulated product (Mudalip et al., 2021; Akbarbaglu et al., 2019; Castro-López et al., 2021; Valková et al., 2022).

While spray drying has been widely employed for encapsulating bioactive compounds, studies optimizing the encapsulation of flavonoids from Tiwai using this method are limited. Most studies have focused on single-variable optimization, which is time-consuming and does not account for potential interactions among multiple factors (Mehmood et al., 2012). In contrast, Response Surface Methodology (RSM), specifically the Box-Behnken design (BBD), is a more efficient approach for optimizing complex processes. BBD allows for the simultaneous assessment of multiple factors and their interactions, making it ideal for optimizing the encapsulation process by adjusting the ratio of Tiwai extract to maltodextrin (ET:MD), chitosan concentration, and inlet temperature. This methodology also helps maximize the range of independent variables and provides valuable data on factors influencing key responses such as encapsulation efficiency (EE), loading capacity (LC), total flavonoid content (TFC), and antioxidant activity (DPPH) (Yolmeh & Jafari, 2017; Raj & Murugesan, 2022).

This study aims to optimize the encapsulation of Tiwai extract using the spray drying method. Specifically, the study focuses on adjusting the ET:MD ratio, chitosan concentration, and inlet temperature to achieve the highest possible EE, LC, and TFC while also assessing antioxidant activity. The encapsulated product will be characterized using FTIR analysis to evaluate changes in functional groups, and flavonoid content will be quantified using HPLC to ensure that the encapsulation process enhances both the stability and bioavailability of the flavonoids. These findings will have potential

applications in functional foods and pharmaceuticals, offering a promising solution to the instability of flavonoids in Tiwai extract.

## 2. Objectives

The objective of this study was to optimize the encapsulation of Tiwai (*Eleutherine americana* L. Merr) extract using Response Surface Methodology (RSM) with the Box-Behnken design (BBD) and to evaluate the stability, potential bioavailability, and antioxidant activity of the encapsulated flavonoids. The study also aimed to assess key parameters, including encapsulation efficiency (EE), loading capacity (LC), total flavonoid content (TFC), and antioxidant activity (DPPH), while characterizing functional groups through FTIR analysis and identifying flavonoids using HPLC for potential applications in functional foods and pharmaceuticals.

## 3. Materials and Methods

### 3.1 Preparation of Tiwai Extract

The seven-month harvested Tiwai onion was washed with running water and sliced using a Tupperware slicer (1 mm thickness). Drying was carried out using a microwave at 100 °C for 30 minutes based on the study by Rachmawati et al. (2024). This method was selected due to its faster drying time compared with conventional oven and freeze-drying methods, and it better preserves the bioactive flavonoids by minimizing heat-induced degradation. Pulverization was carried out using a grinder, and the powdered Tiwai was sieved using an 80-mesh sieve. For extraction, 10 g of Tiwai powder was dissolved in 100 mL of distilled water and macerated for 24 hours. Subsequently, evaporation was performed using an oven at 50 °C until a concentrated extract was obtained. The final extract yield was calculated as a percentage weight-to-weight (w/w) of the dried extract relative to the initial raw material.

### 3.2 Chemicals and Reagents

Tiwai onion was obtained from local farmers in Samarinda, Indonesia. Folin-Ciocalteu reagent was supplied by Fisher (Loughborough, UK). Sodium carbonate, quercetin, potassium ferrocyanide, and acetic acid were obtained from Scharlau (Sentmenat, Spain). Ascorbic acid and 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical were purchased from Sigma (Steinheim, Germany), and iron (II) chloride tetrahydrate was obtained from Panreac Química SAU (Barcelona,

Spain). All other chemicals and reagents used were of analytical grade and commercially available.

### 3.3 Variable Selection and Encapsulation Process

The weight ratio of Tiwai extract to maltodextrin ( $X_1$ ) ranged from 0.1375 to 0.2625 g/g, chitosan concentration ( $X_2$ ) varied from 0.06 to 0.24 g, and the spray-drying inlet temperature ( $X_3$ ) ranged from 70 to 80°C (Table 1). These values were selected based on a previous study of encapsulant product formulations (Liang et al., 2017). The dependent variables measured were encapsulation efficiency (EE), loading capacity (LC), total flavonoid content (TFC), and DPPH radical scavenging activity. The Tiwai extract was mixed with the encapsulating materials in varying ratios and processed at different inlet temperatures as specified in Table 1. The experimental design was generated using Design-Expert software version 13.0 (Stat-Ease, Inc., Minneapolis, MN, USA). The compressor pressure and flow rate in the spray drying device were stabilized at -40 mmHg and 5 to 20 mL per minute, respectively.

### 3.4 Experiment Design

Optimization experiments were conducted using RSM for EE, LC, TFC, and DPPH methods. BBD consisted of 15 experiments including three

center point experiments (Table 1). Encapsulation formula variables included the (ET:MD) ratio ( $X_1$ , g), chitosan concentration ( $X_2$ , g), and inlet temperature (SD) ( $X_3$ , °C). The experimental data were fit with a second-order polynomial model to obtain the regression coefficient ( $\beta_0$ ). The general second-order polynomial model used in response surface analysis is as follows:

$$Y = \beta_0 + \sum_{i=1}^k \beta_i X_i + \sum_{i=1}^k \beta_{ii} X_i^2 + \sum_{i < j}^k \beta_{ij} X_i X_j \quad (1)$$

where Y is the response variable,  $X_i$  and  $X_j$  are the independent variables, n is the number of variables tested, and  $\varepsilon$  represents the error.  $\beta_0$ ,  $\beta_i$ ,  $\beta_{ii}$ , and  $\beta_{ij}$  are the constant, linear, quadratic, and interaction effect coefficients, respectively.

To evaluate the predictive model for the response variables, analysis of variance (ANOVA) was conducted at a 95% confidence level to assess the effect of each factor, namely the Tiwai extract-to-maltodextrin (ET:MD) ratio, chitosan concentration, and spray-drying inlet temperature. The coefficient of determination ( $R^2$ ), p-value of the regression model, and the lack-of-fit (LOF) p-value were examined to determine the adequacy of the regression model. Optimal conditions were identified based on response surface analysis (3D plots).

**Table 1** Experimental factors and measured response values

Run	Independent Variables			Responses			
	Ratio (ET:MD) ( $X_1$ )	Chitosan Concentration ( $X_2$ )	Inlet Temperature (SD) ( $X_3$ )	EE (%)	LC (%)	TFC (mg QE/g)	DPPH ( $\mu\text{g/mL}$ )
1	0.2	0.24	80	65.03	80.79	44.02	63.07
2	0.2	0.15	75	96.85	94.656	65.56	65.17
3	0.1375	0.24	75	53.36	68	36.12	55.62
4	0.1375	0.15	70	72.05	93.46	48.77	152.1
5	0.2625	0.15	80	65.9	94.22	44.61	124.7
6	0.2	0.15	75	95.12	95.185	67.23	63.1
7	0.1375	0.06	75	66.63	96.44	45.1	55.06
8	0.2	0.24	70	60.14	84.17	40.71	52.77
9	0.1375	0.15	80	73.86	88.65	72.54	129.3
10	0.2625	0.06	75	49.29	94.95	33.36	113.1
11	0.2	0.15	75	96.03	93.85	63.53	62.87
12	0.2	0.06	80	71.96	95.71	48.71	46.35
13	0.2	0.06	70	46.53	95.4	31.5	17.37
14	0.2625	0.15	70	69.78	93.15	47.23	74.51
15	0.2625	0.24	75	59.75	86.81	40.45	39.14

*Notes.* Factors: ratio Tiwai extract (ET): maltodextrin (MD) ratio (g), chitosan concentration (g), and inlet temperature spray drying (SD) (°C). The dependent variables measured were encapsulation efficiency (EE), loading capacity (LC), total flavonoid content (TFC), and antioxidant activity (DPPH method)

### 3.5 Validation of Model

The optimized encapsulation conditions, namely the ratio of Tiwai extract to maltodextrin (ET:MD), chitosan concentration, and spray drying inlet temperature, were validated based on the parameters of EE, LC, TFC, and DPPH, which were obtained using RSM. All these variables were re-established in the encapsulation process at optimum conditions. Subsequently, the experimental values were compared with the predicted values of the model to assess the validity of the optimization. The functional groups in the constituent materials and encapsulated Tiwai extract were analyzed using ATR-FTIR, whereas the flavonoid profile was determined using High-Performance Liquid Chromatography (HPLC) under optimum conditions.

#### 3.5.1 Encapsulation Efficiency and Loading Capacity

Flavonoids in the Tiwai extract encapsulant were determined using a spectrophotometric method with an aluminum chloride ( $\text{AlCl}_3$ ) reagent. A total of 50 mg of encapsulant powder was dissolved in 10 mL of 95% ethanol and then sonicated for 30 minutes to extract the encapsulated flavonoids. The sonication solution was centrifuged at 4000 rpm for 15 min, and the supernatant obtained was used as the test sample. A total of 1 mL of supernatant was taken and added to 0.7 mL of distilled water and 0.1 mL of 5%  $\text{NaNO}_2$  solution, and the mixture was allowed to stand for five minutes. Next, 0.1 mL of 10%  $\text{AlCl}_3$  solution was added and allowed to stand for six minutes, followed by the addition of 0.5 mL of 1 M NaOH. The mixture was homogenized and incubated for 10 minutes. The absorbance was measured at a wavelength of 510 nm using a UV-Vis spectrophotometer, with 95% ethanol as a blank. This was followed by calculating flavonoid content using a quercetin standard curve and expressed as milligrams of quercetin equivalent per gram of sample (mg QE/g) (Zou et al., 2004). The EE value was calculated based on the ratio between the amount of encapsulated flavonoids and the initial amount of flavonoids used. Meanwhile, LC was calculated as the percentage of the amount of flavonoids relative to the total encapsulant weight. All tests were performed in triplicate to improve the accuracy of the results.

#### 3.5.2 Analysis of TFC Content

An extract sample (1 mg) was dissolved in 10 mL of 95% ethanol. Subsequently, 0.7 mL of distilled

water and 0.1 mL of 5%  $\text{NaNO}_2$  solution were added. After 5 min, 0.1 mL of 10%  $\text{AlCl}_3$  solution was introduced, followed 6 min later by 0.5 mL of 1 M NaOH. The mixture was then homogenized and incubated for 10 min. Absorbance was recorded at 510 nm, with 1 mL of 95% ethanol used as the blank. The total flavonoid content (TFC) was calculated and expressed as milligrams of quercetin equivalent per gram of dry weight (mg QE/g) (Wunnakup et al., 2024; Zou et al., 2004).

#### 3.5.3 Antioxidant Activity (DPPH)

The DPPH radical capture ability test was conducted based on a modified method from Chew et al. (2009). A total of 1 mL of plant extracts in various concentrations diluted with ethanol was mixed with 1 mL of DPPH solution (0.15 mM in ethanol). As a control, a mixture of 1 mL of DPPH solution with 1 mL of ethanol was prepared. The reaction mixture was shaken until homogeneous, then incubated in the dark at room temperature for 30 minutes. After incubation, the absorbance was measured at a wavelength of 517 nm using an Eppendorf Bio Spectrometer Basic. Ascorbic acid was used as a positive control, and ethanol was used as a blank. The capture ability of DPPH radicals by plant extracts was calculated using the following equation:

$$\% \text{ DPPH capture activity} = \frac{(\text{Abs control} - \text{Abs sample})}{\text{Abs control}} \times 100$$

where Abs control is the absorbance of the mixture of DPPH and ethanol, whereas Abs sample is the absorbance of the mixture of DPPH with the sample. Additionally, three types of controls were prepared to validate the results.

#### 3.5.4 FTIR Analysis

FTIR spectra were scanned using a German FTIR spectrophotometer (Bruker Alpha II). Measurements were taken in the mid-infrared region 4000–600  $\text{cm}^{-1}$  with a resolution of 4  $\text{cm}^{-1}$ . A horizontal attenuated total reflection (ATR) accessory equipped with ZnSe crystals was used for sampling. All FTIR spectra were baseline-corrected using the spectrum of air as the background, with a new reference air spectrum collected after each scan. Spectra were recorded in absorbance mode, with each measurement performed in triplicate. These data were then used to establish correlations between FTIR spectra, antiradical activity, and flavonoid content.

### 3.5.5 HPLC Analysis of Flavonoids under Optimum Conditions

Analysis of flavonoids in Tiwai extract obtained under optimum conditions was carried out for three replicates using a High-Performance Liquid Chromatography (HPLC) HP1200 with a diode array detector (Agilent Technologies, Palo Alto, CA, USA). The separation was performed using an Inertsil ODS-3 column (5.0  $\mu\text{m}$  particle size; 4.6 mm  $\times$  250 mm) equipped with an Inertsil ODS-3 guard column (5.0  $\mu\text{m}$ ; 4.0 mm  $\times$  10 mm) from GL Sciences Inc. (Tokyo, Japan). The elution conditions used a gradient system with solvent A as 2.5% formic acid in water and solvent B as 2.5% formic acid in acetonitrile. The elution gradient profile was arranged as follows: 0 min (5% B), 13 min (11% B), 16 min (13% B), 20 min (14% B), 22 min (15% B), 25 min (20% B), 28 min (25% B), 30 min (30% B), and back to 5% B at the 40th minute. The total analysis time was 40 minutes, with a solvent flow rate of 1 mL/minute and an injection volume of 25  $\mu\text{L}$ . Identification of flavonoids was carried out by comparing the retention time of sample chromatogram peaks against reference standards (Suwanpitak et al., 2025). The concentration of individual phenolic compounds was calculated using a calibration curve that had been previously made (Hongsing et al., 2024). The results were expressed in units of micrograms per gram ( $\mu\text{g/g}$ ) of sample.

## 4. Results and Discussions

### 4.1 Response Surface Model and Analysis

The determination of variable levels in the encapsulant formula, including the ratio of Tiwai extract to maltodextrin (ET:MD), chitosan concentration, and spray-drying inlet temperature, was based on preliminary investigations. These variables were incorporated into the experimental design to evaluate their impact on encapsulated characteristics. The selected variables, ET:MD ratio, chitosan concentration, and spray-drying inlet temperature were chosen due to their significant influence on key encapsulation parameters such as encapsulation efficiency (EE), loading capacity (LC), total flavonoid content (TFC), and antioxidant activity (DPPH). These variables have been extensively studied in previous literature and are well-known for their effects on the stability and bioavailability of active compounds in encapsulated products (Hong et al., 2016; Yoksan et al., 2010). The ET:MD ratio affects the interaction between the active ingredient and the encapsulant, while chitosan concentration plays a

crucial role in determining the protective capability of the encapsulation matrix. The spray-drying temperature is an essential parameter for achieving stable encapsulants. For optimization and statistical analysis, Box-Behnken design (BBD) based on Response Surface Methodology (RSM) was employed. This method facilitates regression analysis and response surface visualization, enabling the identification of optimal conditions by analyzing the interactions between the selected variables.

### 4.2 Fitting the Model

The encapsulation formula was optimized using a second-order polynomial model. The model exhibited high statistical significance and an excellent fit to the experimental data for EE, LC, TFC, and DPPH, with corresponding  $R^2$  values of 0.9572, 0.9515, 0.9682, and 0.8051, respectively (Table 2). The regression coefficients for the dependent variables were obtained by multiple linear regression as shown in Table 2. Based on the results, the linear effect of (ET:MD) ratio ( $X_1$ ) was significant for LC and TFC, chitosan concentration ( $X_2$ ) for LC, and SD inlet temperature for TFC. The interaction effect of  $X_{12}$  (ET:MD ratio and chitosan concentration) was found for LC, whereas  $X_{13}$  (ET:MD ratio and SD inlet temperature) was only found for TFC. The quadratic effect of  $X_1$  (ET:MD ratio) had significant negative effects on EE and TFC, but positive effects on DPPH. The square effect of  $X_2$  (chitosan concentration) produced significant negative effects on EE, LC, and TFC. Additionally, the square effect of  $X_3$  (SD inlet temperature) also produced negative effects on EE.

The ANOVA results for EE, LC, and TFC indicated that the three model parameters effectively explained the variation in the experimental data for these response variables. This was evidenced by the significant F-values for the models (Table 2) and the non-significant lack-of-fit tests, confirming that the models adequately represented the experimental data. In contrast, the DPPH variable showed a non-significant model effect, suggesting that the experimental data were insufficient to accurately describe this response.

### 4.3 Effect of Encapsulation Formula Variables on EE

The model showed high significance values ( $p < 0.01$ ) with the experimental data. ANOVA showed a negative ( $p = 0.01$ ) linear effect ( $X_1$ ), quadratic effect of  $X_1$  ( $p < 0.01$ ),  $X_2$  ( $p < 0.001$ ) and  $X_3$  ( $p < 0.05$ ) on the EE response, as presented in Table 2.

There was no significant interaction effect between  $X_1$ ,  $X_2$ , and  $X_3$  on the experimental data. The insignificant discrepancy value ( $F = 0.1395$ ;  $p > 0.05$ ) showed that the model fit the effects of the selected variables on this response with good prediction ( $R^2 = 0.9572$ ) and was nonsignificant compared to pure error.

EE values obtained in the study were similar to those of Ding et al. (2019), who also used maltodextrin as a coating material. The results of regression analysis on each quadratic variable of the ratio (ET:MD), maltodextrin concentration, and SD inlet temperature had a significant effect on EE. This showed that the relationship between the variables and EE response was non-linear. More importantly, an increase or decrease in one of the variables does not significantly cause a linear or unidirectional change in the response (Kalam et al., 2016). Positive values determine synergistic effects on optimization, while negative values direct antagonistic effects between factors and responses (Hao et al., 2011).

The relationship between EE and process variables is shown in Figure 1a. The highest EE occurred in the range of 0.1375g to 0.200 g. Each variable and its interaction did not have a significant effect on EE in determining the Tiwai extract encapsulant formula, but a substantial impact was observed for the quadratic effect. This suggested that EE did not increase or decrease proportionally with a rise in  $X_1$ ,  $X_2$ , and  $X_3$ , but through a more complex pattern (Rahman et al., 2010).

#### 4.4 Effect of Encapsulation Formula Variables on LC

The model showed significant values ( $p < 0.05$ ) in the experimental data. ANOVA showed significant effects of  $X_1$  ( $p < 0.05$ ) and  $X_2$  ( $p < 0.001$ ), the interaction between variables  $X_1$  ( $p < 0.05$ ), and a negative quadratic effect of  $X_2$  ( $p < 0.05$ ) on the LC response, as shown in Table 2. The nonsignificant discrepancy value ( $F = 1.75$ ;  $p > 0.05$ ) suggested that the model adequately fit the spatial influence of the variables on this response, with good prediction ( $R^2 = 0.9515$ ). The LC values obtained were more than 80% similar to the results reported by Permanadewi et al. (2022), who used spray-drying as the encapsulation method. Additionally, regression analysis results showed higher LC at the ratio of ET:MD, as well as greater chitosan concentration up to a certain limit. The ratio of encapsulant and coating materials used played an important role in LC obtained in the encapsulation of rosmarin essential oil (Turasan et al.,

2015). The relationship between LC and process variables is shown in Figure 1b, where the highest LC values range from 0.12 to 0.2375% and 0.1875 to 0.2375%.

The encapsulant material ratio (ET:MD) and chitosan concentration play a crucial role in the LC of the product (Yoksan et al., 2010). The balance between extract (ET) as the active ingredient and maltodextrin (MD), including chitosan as the coating material, can significantly affect the ability of the encapsulant to effectively encapsulate and protect the active ingredient. This suggests that a greater amount of extract may increase LC. However, increasing the amount of extract beyond a certain point could weaken the structural integrity of the wrapping matrix due to its inability to stabilize the core material properly. On the other hand, increasing the content of maltodextrin and chitosan generally improves the structural stability of the matrix but reduces the proportion of extracts as active ingredients. Several studies have reported that the ratio of encapsulant material (ET) and chitosan concentration determines both encapsulation efficiency (EE) and LC. This is attributed to the significant effect on structural stability, protection of the core material, and the controlled release characteristics of the encapsulated substance (Ren et al., 2019; Soltanzadeh et al., 2021; Woranuch & Yoksan, 2013).

#### 4.5 Effect of Encapsulation Formula Variables on TFC

ANOVA revealed significant linear effects and negative quadratic effects of  $X_1$ ,  $X_3$ ,  $X_1^2$ , and  $X_2^2$  on TFC values, with  $p < 0.05$  for all variables except  $X_2^2$  ( $p < 0.001$ ) (Table 2). Additionally, the interaction between the ET:MD ratio and spray-drying inlet temperature was highly significant ( $p < 0.001$ ). As shown in Figure 1c, the ET:MD ratio, chitosan concentration, and inlet spray-drying temperature for TFC reached the highest point before decreasing. This trend was consistent with the EE results, where the parameters showed a strong positive correlation ( $R^2 = 0.89$ ,  $p < 0.001$ ) (Table 3). The highest flavonoid content was observed at an ET:MD ratio of 0.1875–0.2125 g (Figure 1c), chitosan concentration of 0.155–0.18 g, and spray-drying inlet temperature of 74–76°C (Figure 1d). The positive correlation between EE and TFC showed a unidirectional relationship between the two variables. This relationship could be due to the better ability of the encapsulation matrix to protect or

maintain flavonoids during the encapsulation process (Ydjedd et al., 2017).

#### 4.6 Effect of Encapsulation Formula Variables on DPPH

ANOVA revealed significant linear effects and negative quadratic effects of  $X_1$ ,  $X_3$ ,  $X_1^2$ , and  $X_2^2$  on DPPH values, with  $p < 0.05$  for all variables except  $X_2^2$  ( $p < 0.001$ ) (Table 2). Additionally, the interaction between the ET:MD ratio and spray-drying inlet temperature was highly significant ( $p < 0.001$ ). As shown in Figure 1d, the DPPH value increased as the ET:MD ratio decreased, indicating a reduction in antioxidant capacity, as reflected by higher  $IC_{50}$  values (Patthamasopasakul et al., 2024). This suggests that a lower ET:MD ratio leads to reduced antioxidant activity, possibly due to a decrease in the effective encapsulation of flavonoids and other active compounds. The DPPH response, measured as the inhibitor concentration ( $IC_{50}$ ), showed a quadratic effect of the ET:MD ratio ( $X_1^2$ ) being significant ( $p < 0.05$ ) (Table 2), indicating a non-linear relationship between the ET:MD ratio and antioxidant capacity. This result suggests that a balance in the ET:MD ratio is crucial for optimizing antioxidant activity, where


very high or low ratios may lead to less effective encapsulation and reduced antioxidant performance. However, the model did not fully describe the observed behavior, as indicated by the lack-of-fit F-value (810.43, highly significant). This suggests that the current model does not fully capture the complexity of the system, and additional variables or a more refined model might be necessary to better predict antioxidant activity. Factors not considered by the model could include the degradation of flavonoids during the encapsulation process, interactions between flavonoids and other matrix components, and non-flavonoid antioxidant contributions that may influence the DPPH response.

The correlation of each variable can be observed in Table 3. The low correlation between TFC and DPPH was due to the contribution of other antioxidant compounds, variations in the type and structure of flavonoids, differences in measurement methods, interactions among compounds in the encapsulation matrix, and the possibility of flavonoid degradation during the encapsulation process. Similarly, Sulaiman et al. (2011) reported a low correlation between TFC and DPPH.

**Table 2** Regression coefficient ( $\beta$ ), coefficient of determination ( $R^2$ ), and F-test values of the predicted second order polynomial models for EE, LC, TFC, and DPPH

	Regression Coefficients ( $\beta$ )			
	EE	LC	TFC	DPPH
Intercept				
$X_0$	96.68	96.33	65.44	63.71
Linear				
$X_1$	-2.65	2.82*	-4.61*	-5.08
$X_2$	0.4837	-7.84***	0.3287	-2.66
$X_3$	3.53	-0.8513	5.21*	8.33
Cross product				
$X_{12}$	5.93	5.08*	4.02	-18.63
$X_{13}$	-1.42	1.47	-6.60*	18.25
$X_{23}$	-5.13	-0.9225	-3.48	-4.67
Quadratic				
$X_1^2$	-14.97**	-3.22	-7.32*	38.64*
$X_2^2$	-24.45***	-6.57*	-19.37***	-36.62
$X_3^2$	-11.31*	-0.7483	-4.84	17.80
$R^2$	0.9572	0.9515	0.9682	0.8051
F value (model)	12.44**	10.91**	16.91*	2.29
F value (lack of fit)	0.1395	1.75	6.41	810.43*

Notes. Note.  $X_1$  = ET:MD ratio (g);  $X_2$  = chitosan concentration (g);  $X_3$  = spray-drying inlet temperature ( $^{\circ}C$ );  $X_1^2$ ,  $X_2^2$ ,  $X_3^2$  = quadratic effects;  $X_{12}$ ,  $X_{13}$ ,  $X_{23}$  = interaction effects; EE = encapsulation efficiency (%); LC = loading capacity (%); TFC = total flavonoid content (mg QE/g); DPPH = 2,2-diphenyl-1-picrylhydrazyl radical scavenging activity ( $\mu g/mL$ );  $R^2$  = coefficient of determination. \*  $p < 0.05$ ; \*\*  $p < 0.01$ ; \*\*\*  $p < 0.001$ .



**Figure 1** Effect of variables on the response of a) Encapsulation efficiency, b) Loading capacity, c) Total flavonoid content (TFC) mg QE/g, and d) Antioxidant activity (DPPH) µg/mL Three-dimensional (3D) response surface plots

**Table 3** Pearson correlations calculated for response variables

	EE	LC	TFC	DPPH
EE	1.00	0.35	0.89**	0.09
LC		1.00	0.27	0.15
TFC			1.00	0.26
DPPH				1.00

Notes. Level of significance \*  $p < 0.05$ ; \*\*  $p < 0.01$ ; EE = Encapsulation efficiency; LC = Loading Capacity; TFC = Total Flavonoid Content (mg QE/g); DPPH = 2,2- diphenyl-1-picrylhydrazyl radical scavenging ability (µg/mL)

**Table 4** Experimental data validating predicted values for optimal encapsulation conditions of Tiwai extract

Dependent Variable	Predicted Value	Experiment Value	%CV
EE (%)	95.97	93.24	2.86
LC (%)	97.53	95.76	1.81
TFC (mg QE/g)	65.55	69.23	5.62
DPPH ( $\mu\text{g}/\text{mL}$ )	63.55	60.01	5.56

*Note.* EE = encapsulation efficiency; LC = loading capacity; TFC = total flavonoid content; DPPH = 2,2-diphenyl-1-picrylhydrazyl radical scavenging activity; %CV = percentage coefficient of variation. Predicted values were generated using Design-Expert software version 13.0 based on the optimized Box–Behnken design.

#### 4.7 Encapsulation Formula Parameter Optimization and Model Validation

Optimum conditions were identified by maximizing the overall desirability using Design-Expert software version 13.0 (Stat-Ease, Inc., Minneapolis, MN, USA). The optimization aimed to achieve the highest possible EE, LC, and TFC values, while minimizing DPPH. The resulting optimal formulation parameters were an ET:MD ratio of 0.195 g, chitosan concentration of 0.130 g, and spray-drying inlet temperature of 75°C, yielding a desirability score of 0.805. These conditions were applied in the encapsulation process, and the responses were determined and validated following standard procedures. The experimental results closely matched the predicted values, with coefficients of variation (CV) ranging from 1.81 to 5.62 (Table 4). This parameter describes the extent to which the data are scattered. Under optimal conditions for the Tiwai extract encapsulant formula, additional measurements of functional groups were made for comparison with the raw material. The quercetin compound content of the final product was also identified using the HPLC method.

#### 4.8 Scanning of FTIR Spectra

Figure 2 shows the FTIR spectra of the encapsulant samples of Tiwai extract (TEE), TE, with Tiwai extract and a mixture of maltodextrin and chitosan (TE+CM) scanned in the mid-infrared region corresponding to wave numbers 4000–650  $\text{cm}^{-1}$ . Each peak in FTIR spectra corresponds to the stretching and bending vibrations of functional groups that absorb radiation, causing vibrational transitions between bonds. The shape of the FTIR spectra in the three samples shows the presence of functional groups that differ in absolute intensity. This is because the FTIR spectra describe the interaction of infrared light with molecules in the sample, leading to a characteristic absorption pattern (Chen et al., 2012; Siddique, 2024).


The interpretation of the FTIR spectra in term of wave numbers, functional groups, and transmittance is shown in Table 5. Wave number (a) 3276.34  $\text{cm}^{-1}$  shows

a low stretching of –OH (hydroxyl bond) in TEE. This may be caused by the heating process during spray drying leading to a decrease in free hydroxyl groups or converting into ether or ester groups (Akbarbaglu et al., 2019). Flavonoids with a carbonyl group (C=O) at the ketone position in ring C (e.g., flavones and flavonols such as quercetin) and aromatic compounds such as rings A and B show C=C vibration at 1638.78  $\text{cm}^{-1}$  peak (Chen et al., 2022; Kolekar et al., 2019). Moreover, the lower transmittance intensity in TEE (63.6%) compared with TE+CM (70.6%) indicates that the flavonoid content in TEE is higher than TE+CM. This occurs because of free or active carbonyls (Horincar et al., 2019), including the coating material in TE+CM that can reduce the availability of flavonoid aromatic groups to absorb infrared light due to imperfect homogenization (Rahmadi et al., 2022).

The peak at 1413.8275  $\text{cm}^{-1}$  is associated with C–H deformation vibrations derived from aliphatic or aromatic structures and stretching vibrations of C–O groups in compounds with carboxylic (–COO–) or ester groups (Yang et al., 2016a; Yang et al., 2016b). However, this peak is not detected in the encapsulant of Tiwai extract due to the reaction of active groups with the coating material (maltodextrin and chitosan) during the encapsulation process (Rahmadi et al., 2022). The peak at 1357–1361.10  $\text{cm}^{-1}$ , attributed to C–H bending (aromatic/aliphatic), such as –CH<sub>2</sub> and –CH<sub>3</sub> (Russo et al., 2014), the transmittance of 90.4% in TEE shows low absorption. This indicates that the content of aromatic/aliphatic C–H groups is less active compared with TE and TE+CM. Furthermore, C–O/C–N and C–O stretching functional groups at peaks 1149.93, 1017.69, and 926.69  $\text{cm}^{-1}$  are undetectable or minor in TE. These functional groups include C–O stretching derived from ether and ester bonds, or carbohydrate compounds such as maltodextrin. C–N stretching is commonly found in amine or amide compounds such as chitosan, and C–O Stretching (Glycosidic Linkage) is derived from glycosidic bonds in polysaccharides (maltodextrin) or glycosides (Fornaro et al., 2016). (g)

The peak at 1076.7795  $\text{cm}^{-1}$  is related to C–O–C stretching vibration, which likely corresponds to glycosidic linkages or ether bonds found in the encapsulant material, such as maltodextrin and

chitosan. The vibrations observed in the FTIR spectra support the presence of flavonoids in the Tiwai extract encapsulant, which are responsible for the antioxidant activity.



**Figure 2.** FTIR spectra of TEE (Tiwai Extract Encapsulant), TE (Tiwai Extract), TE+CM (Tiwai Extract + Coating Material) scanned at wave numbers 4000–650  $\text{cm}^{-1}$  using the damped total reflectance (ATR) mode

**Table 5** FTIR functional group comparison

Code	Wave Number ( $\text{cm}^{-1}$ )	Functional Group	TEE (%T)	TE (%T)	TE+CM (%T)
a	3276.34	O–H stretching (hydroxyl)	870	481	492
b	2910 – 2932	C–H stretching (aliphatic/aromatic)	910	956	842
c	1638.78	C=O stretching (carbonyl) / C=C (aromatic)	636	657	706
d	1413.82	C–H bending / C–O (carboxylate)	0	754	783
e	1357 – 1361.10	C–H bending (aromatic/aliphatic)	904	751	779
f	1149.93	C–O / C–N stretching	847	0	633
g	1076.7795	C–O–C stretching	768	756	520
h	1017.6902	C–O stretching (ether) / C–N (amine)	928	819	271
i	926.6942	C–O stretching (glycosidic linkage)	873	0	715

**Table 6** Quercetin standard and encapsulated concentration of Tiwai extract

Concentration	Real Time Min (25)	Area mAU* min (25)	Height mAU	Amount ( $\mu\text{g/g}$ )
Quercetin				
100	18.47	6.85	25.65	133.94
200	18.46	19.03	72.54	224.15
500	18.41	42.82	177.13	400.26
Tiwai extract encapsulant (TEE)				
	18.37	8.79	38.35	148.30

#### 4.9 HPLC Analysis of Flavonoids

Qualitative and quantitative analysis of phenolic compounds from the extracts obtained under optimum conditions (30°C, 25 min) using the HPLC method showed that the quercetin content in the Tiwai extract encapsulant was 148.30 µg/g. This value (0.148 mg/g when converted to consistent units) was substantially lower than the TFC of 69.23 mg QE/g measured by the AlCl<sub>3</sub> colorimetric test, reflecting the method's broader reactivity with non-flavonoid antioxidant compounds. The difference in results can be attributed to variation in the basic principles of the two methods as the AlCl<sub>3</sub> method not only reacted with flavonoids, but also responded to other antioxidant compounds such as proteins, carbohydrates, and unsaturated fatty acids, providing a higher estimate of flavonoids (Sultana et al., 2024; Santos-Sánchez et al., 2019; Gulcin, 2020; Munteanu & Apetrei, 2021). In contrast, the HPLC method is more selective and specific to certain flavonoids, such as quercetin, and thus provides more accurate values, although lower in quantity (Deshmukh & Patilm, 2024; Jain & Shaikh, 2016; Nguyen et al., 2023). Similar results were also reported by Sultana et al. (2024), who found that AlCl<sub>3</sub>-based derivatization in HPTLC analysis led to higher flavonoid estimates compared with chromatographic quantification due to the broad reactivity of AlCl<sub>3</sub> with non-flavonoid antioxidant molecules. This emphasizes the importance of method selection in determining flavonoid content, particularly when comparing total flavonoids and individual compounds such as quercetin.

The results were in line with the FTIR analysis, which showed significant changes in key functional groups in the flavonoid structure, such as a decrease in the intensity of free hydroxyl groups (–OH), carbonyl groups (C=O), and aromatic bonds (C=C) in the encapsulant sample. This indicated a chemical interaction between the active compound (such as quercetin) and the coating material (maltodextrin and chitosan) during the encapsulation process, which could inhibit quercetin in its free form. The absence of the C–H group at wave number 1413.8275 cm<sup>-1</sup> and the appearance of new functional groups such as C–O/C–N stretching further confirmed structural changes in the encapsulant attributable to molecular interactions during encapsulation.

#### 5. Conclusion

In conclusion, formulation and optimization of Tiwai extract encapsulant using the spray drying method were successfully carried out using BBD-RSM. The ET:MD ratio, chitosan concentration, and inlet temperature were shown to have significant effects on EE, LC, TFC, and DPPH. Optimum condition produced encapsulant with high efficiency (EE 93.24%; LC 95.76%), adequate flavonoid content (TFC 69.23 mg QE/g), and antioxidant activity (DPPH 60.01 µg/mL). In line with the results, FTIR analysis showed changes in functional group structure due to the interaction of active compound with coating matrix, which strengthened the evidence of successful encapsulation. Quercetin content measured by HPLC (148.30 µg/g) showed a selective profile of active compounds that was not detected by the conventional colorimetric method.

#### 6. Abbreviations

Abbreviation	Full term
AlCl <sub>3</sub>	Aluminum chloride
ANOVA	Analysis of variance
ATR	Attenuated total reflectance
BBD	Box–Behnken design
DPPH	2,2-Diphenyl-1-picrylhydrazyl
EE	Encapsulation efficiency
ET	Tiwai extract
ET:MD	Ratio of Tiwai extract to maltodextrin
FTIR	Fourier transform infrared spectroscopy
HPLC	High-performance liquid chromatography
LC	Loading capacity
MD	Maltodextrin
RSM	Response surface methodology
SD	Spray drying
TEE	Tiwai extract encapsulant
TE	Tiwai extract
TE+CM	Tiwai extract with coating material
TFC	Total flavonoid content
QE	Quercetin equivalent

Abbreviation	Full term
IC <sub>50</sub>	Half maximal inhibitory concentration
CV	Coefficient of variation

## 7. Acknowledgements

The author is grateful to the Directorate General of Higher Education, Ministry of Education, Culture, Research and Technology, Indonesia, for the funding provided through the Doctoral Research Scheme 2023 with contract number: 492/UN17.L1/HK/2023.

## 8. CRediT Statement

**Maulida Rachmawati:** Investigation, Data Curation, Formal Analysis, Writing – Original Draft.

**Vivie Septianasari:** Methodology, Validation, Writing – Review & Editing.

**Sarah Maulida:** Investigation, Data Curation.

**Rimbawan Apriadi:** Formal Analysis, Visualization. Hadi Suprpto: Resources, Validation.

**Wiwit Murdianto:** Supervision, Writing – Review & Editing.

**Bernatal Saragih:** Conceptualization, Methodology, Supervision, Project Administration, Funding Acquisition, Writing – Review & Editing.

## 9. References

- Adesina, S., Iwalewa, E., & Johnny, I. (2016). Tetrapleura tetraptera Taub-ethnopharmacology, chemistry, medicinal and nutritional values-a review. *British Journal of Pharmaceutical Research*, 12(3), 1-22. <https://doi.org/10.9734/BJPR/2016/26554>
- Akbarbaglu, Z., Jafari, S. M., Sarabandi, K., Mohammadi, M., Heshmati, M. K., & Pezeshki, A. (2019). Influence of spray drying encapsulation on the retention of antioxidant properties and microstructure of flaxseed protein hydrolysates. *Colloids and Surfaces B: Biointerfaces*, 178, 421-429. <https://doi.org/10.1016/j.colsurfb.2019.03.038>
- Castro-López, C., Espinoza-González, C., Ramos-González, R., Boone-Villa, V. D., Aguilar-González, M. A., Martínez-Ávila, G. C., ... & Ventura-Sobrevilla, J. M. (2021). Spray-drying encapsulation of microwave-assisted extracted polyphenols from *Moringa oleifera*: Influence of tragacanth, locust bean, and carboxymethyl-cellulose formulations. *Food Research International*, 144, Article 110291. <https://doi.org/10.1016/j.foodres.2021.110291>
- Chen, P., Cui, H., Feng, L., Yu, J., Hayat, K., Jia, C., ... & Ho, C. T. (2022). Effect of the C-ring structure of flavonoids on the yield of adducts formed by the linkage of the active site at the A-ring and Amadori rearrangement products during the Maillard intermediate preparation. *Journal of Agricultural and Food Chemistry*, 70(10), 3280-3288. <https://doi.org/10.1021/acs.jafc.1c07521>
- Chen, Y., Mastalerz, M., & Schimmelmann, A. (2012). Characterization of chemical functional groups in macerals across different coal ranks via micro-FTIR spectroscopy. *International Journal of Coal Geology*, 104, 22-33. <https://doi.org/10.1016/j.coal.2012.09.001>
- Chew, Y. L., Goh, J. K., & Lim, Y. Y. (2009). Assessment of in vitro antioxidant capacity and polyphenolic composition of selected medicinal herbs from Leguminosae family in Peninsular Malaysia. *Food Chemistry*, 116(1), 13-18. <https://doi.org/10.1016/j.foodchem.2009.01.091>
- Cocuron, J. C., Casas, M. I., Yang, F., Grotewold, E., & Alonso, A. P. (2019). Beyond the wall: High-throughput quantification of plant soluble and cell-wall bound phenolics by liquid chromatography tandem mass spectrometry. *Journal of Chromatography A*, 1589, 93-104. <https://doi.org/10.1016/j.chroma.2018.12.059>
- Deshmukh, M. D., & Patil, M. P. (2024). The quality by design approach for analytical method development and validation of the RP-HPLC method for estimation of quercetin in pure, marketed, and cream formulation. *Biosciences Biotechnology Research Asia*, 21(1), Article 273. <https://doi.org/10.13005/bbra/3224>
- Ding, J., Xu, Z., Qi, B., Cui, S., Wang, T., Jiang, L., ... & Sui, X. (2019). Fabrication and characterization of soybean oil bodies encapsulated in maltodextrin and chitosan-EGCG conjugates: An in vitro digestibility study. *Food Hydrocolloids*, 94, 519-527. <https://doi.org/10.1016/j.foodhyd.2019.04.001>
- Fornaro, T., Biczysko, M., Bloino, J., & Barone, V. (2016). Reliable vibrational wavenumbers for C [double bond, length as m-dash] O and N-H stretchings of isolated and hydrogen-bonded nucleic acid bases. *Physical Chemistry Chemical Physics*, 18(12), 8479-8490. <https://doi.org/10.1039/c5cp07386c>

- Gulcin, İ. (2020). Antioxidants and antioxidant methods: An updated overview. *Archives of Toxicology*, 94(3), 651-715.  
<https://doi.org/10.1007/s00204-020-02689-3>
- Hao, J., Fang, X., Zhou, Y., Wang, J., Guo, F., Li, F., & Peng, X. (2011). Development and optimization of solid lipid nanoparticle formulation for ophthalmic delivery of chloramphenicol using a Box-Behnken design. *International Journal of Nanomedicine*, 6, 683-692.  
<https://doi.org/10.2147/IJN.S17386>
- Hong, S., Shen, S., Tan, D. C. T., Ng, W. K., Liu, X., Chia, L. S., ... & Gokhale, R. (2016). High drug load, stable, manufacturable and bioavailable fenofibrate formulations in mesoporous silica: A comparison of spray drying versus solvent impregnation methods. *Drug Delivery*, 23(1), 316-327.  
<https://doi.org/10.3109/10717544.2014.913323>
- Hongsing, P., Kongart, C., Nuiden, N., Wannigama, D. L., & Phairoh, K. (2024). Quantitative analysis of swertiamarin content from *Fagraea fragrans* leaf extract using HPLC technique and its correlation to antibacterial activity. *Journal of Current Science and Technology*, 14(2), Article 34.  
<https://doi.org/10.59796/jcst.V14N2.2024.341>
- Horincar, G., Aprodu, I., Barbu, V., Râpeanu, G., Bahrim, G. E., & Stănciuc, N. (2019). Interactions of flavonoids from yellow onion skins with whey proteins: Mechanisms of binding and microencapsulation with different combinations of polymers. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 215, 158-167.  
<https://doi.org/10.1016/j.saa.2019.02.100>
- Jain, V. A. N. D. A. N. A., & Shaikh, M. S. (2016). Simultaneous RP-HPLC analysis of quercetin and kaempferol in different plant parts of *Cissus quadrangularis*. *International Journal of Pharmacy and Pharmaceutical Sciences*, 8, 138-42.
- Kalam, M. A., Khan, A. A., Khan, S., Almalik, A., & Alshamsan, A. (2016). Optimizing indomethacin-loaded chitosan nanoparticle size, encapsulation, and release using Box-Behnken experimental design. *International Journal of Biological Macromolecules*, 87, 329-340.  
<https://doi.org/10.1016/j.ijbiomac.2016.02.033>
- Kareem, O., Ali, T., Dar, L. A., Mir, S. A., Rashid, R., Nazli, N., ... & Bader, G. N. (2022). Positive health benefits of saponins from edible legumes: Phytochemistry and pharmacology. *Edible Plants in Health and Diseases: Volume II: Phytochemical and Pharmacological Properties*, 279-298.  
[https://doi.org/10.1007/978-981-16-4959-2\\_8](https://doi.org/10.1007/978-981-16-4959-2_8)
- Kolekar, C., Patil, S. P., & Kumbhar, S. T. (2019). Synthesis and human RBC membrane stabilization activity of substituted 3-benzoyl flavone. *Asian Journal of Pharmacy and Pharmacology*, 5(4), 673-677.  
<https://doi.org/10.31024/ajpp.2019.5.4.4>
- Liang, J., Yan, H., Puligundla, P., Gao, X., Zhou, Y., & Wan, X. (2017). Applications of chitosan nanoparticles to enhance absorption and bioavailability of tea polyphenols: A review. *Food Hydrocolloids*, 69, 286-292.  
<https://doi.org/10.1016/j.foodhyd.2017.01.041>
- Mehmood, T., Liland, K. H., Snipen, L., & Sæbø, S. (2012). A review of variable selection methods in partial least squares regression. *Chemometrics and Intelligent Laboratory Systems*, 118, 62-69.  
<https://doi.org/10.1016/j.chemolab.2012.07.010>
- Mudalip, S. A., Khatiman, M. N., Hashim, N. A., Man, R. C., & Arshad, Z. I. M. (2021). A short review on encapsulation of bioactive compounds using different drying techniques. *Materials Today: Proceedings*, 42(1), 288-296.  
<https://doi.org/10.1016/j.matpr.2021.01.543>
- Munteanu, I. G., & Apetrei, C. (2021). Analytical methods used in determining antioxidant activity: A review. *International Journal of Molecular Sciences*, 22(7), Article 3380.  
<https://doi.org/10.3390/ijms22073380>
- Nguyen, M. H., Ha, D. L., Do, B. M., Chau, N. T. N., Tran, T. H., Le, N. T. H., & Le, M. T. (2023). RP-HPLC-based flavonoid profiling accompanied with multivariate analysis: an efficient approach for quality assessment of *Houttuynia cordata* Thunb leaves and their commercial products. *Molecules*, 28(17), Article 6378.  
<https://doi.org/10.3390/molecules28176378>
- Pathamasopasakul, R., Songsak, T., Kunaratnpruk, S., & Sucontphunt, A. (2024). Comparative study: Extraction conditions and antioxidant and antibacterial activities of *Gracilaria fisheri*. *Journal of Current Science and*

- Technology*, 14(3), Article 52.  
<https://doi.org/10.59796/jcst.V14N3.2024.52>
- Permanadewi, I., Kumoro, A. C., Wardhani, D. H., & Aryanti, N. (2022). Effect of viscosity on iron encapsulation using alginate as a carrying agent in a controlled spray drying process. *Food Research*, 6(5), 56-67.  
[https://doi.org/10.26656/fr.2017.6\(5\).613](https://doi.org/10.26656/fr.2017.6(5).613)
- Rachmawati, M., Murdianto, W., Suprpto, H., Apriadi, R., Ramadhani, N. E., Apriliato, D. K., ... & Saragih, B. (2024). Effect of drying methods and harvest age on the quality of Sliced Tiwai Onion (*Eleutherine americana* Merr) as A Potential Functional Food. *Trends in Sciences*, 21(12), 8401-8401.  
<https://doi.org/10.48048/tis.2024.8401>
- Rahmadi, A., Nurjannah, S., Andriyani, Y., Banin, M. M., Rohmah, M., Amaliah, N., ... & Emmawati, A. (2022). Proximate analysis of the high phytochemical activity of encapsulated Mandai cempedak (*Artocarpus champeden*) vinegar prepared with maltodextrin and chitosan as wall materials. *F1000Research*, 11, Article 865.  
<https://doi.org/10.12688/f1000research.109612.1>
- Rahman, Z., Zidan, A. S., Habib, M. J., & Khan, M. A. (2010). Understanding the quality of protein loaded PLGA nanoparticles variability by Plackett–Burman design. *International Journal of Pharmaceutics*, 389(1-2), 186-194.  
<https://doi.org/10.1016/j.ijpharm.2009.12.040>
- Raj, R. A., & Murugesan, S. (2022). Optimization of dielectric properties of pongamia pinnata methyl ester for power transformers using response surface methodology. *IEEE Transactions on Dielectrics and Electrical Insulation*, 29(5), 1931-1939.  
<https://doi.org/10.1109/TDEI.2022.3190257>
- Ren, X., Hou, T., Liang, Q., Zhang, X., Hu, D., Xu, B., ... & Ma, H. (2019). Effects of frequency ultrasound on the properties of zein-chitosan complex coacervation for resveratrol encapsulation. *Food Chemistry*, 279, 223-230.  
<https://doi.org/10.1016/j.foodchem.2018.11.025>
- Russo, C., Stanzione, F., Tregrossi, A., & Ciajolo, A. (2014). Infrared spectroscopy of some carbon-based materials relevant in combustion: Qualitative and quantitative analysis of hydrogen. *Carbon*, 74, 127-138.  
<https://doi.org/10.1016/j.carbon.2014.03.014>
- Santos-Sánchez, N. F., Salas-Coronado, R., Villanueva-Cañongo, C., & Hernández-Carlos, B. (2019). *Antioxidant Compounds and Their Antioxidant Mechanism*. IntechOpen.  
<https://doi.org/10.5772/intechopen.85270>
- Shahidi, F., & Yeo, J. (2016). Insoluble-bound phenolics in food. *Molecules*, 21(9), Article 1216.  
<https://doi.org/10.3390/molecules21091216>
- Siddique, I. (2024). Exploring functional groups and molecular structures: A comprehensive analysis using FTIR spectroscopy. *Chemistry Research Journal*, 9(2), 70-76.  
<https://doi.org/10.2139/ssrn.4886526>
- Soltanzadeh, M., Peighambaroust, S. H., Ghanbarzadeh, B., Mohammadi, M., & Lorenzo, J. M. (2021). Chitosan nanoparticles as a promising nanomaterial for encapsulation of pomegranate (*Punica granatum* L.) peel extract as a natural source of antioxidants. *Nanomaterials*, 11(6), Article 1439.  
<https://doi.org/10.3390/nano11061439>
- Sulaiman, S. F., Yusoff, N. A. M., Eldeen, I. M., Seow, E. M., Sajak, A. A. B., & Ooi, K. L. (2011). Correlation between total phenolic and mineral contents with antioxidant activity of eight Malaysian bananas (*Musa* sp.). *Journal of Food Composition and Analysis*, 24(1), 1-10. <https://doi.org/10.1016/j.jfca.2010.04.005>
- Sultana, S., Hossain, M. L., Sostaric, T., Lim, L. Y., Foster, K. J., & Locher, C. (2024). Investigating flavonoids by HPTLC analysis using aluminium chloride as derivatization reagent. *Molecules*, 29(21), Article 5161.
- Suwanpitak, K., Sangnim, T., Sriamornsak, P., Puri, V., Sharma, A., & Huanbutta, K. (2025). Development and validation of a reliable reverse-phase high-performance liquid chromatography method for quantifying triterpenes in *Centella asiatica*: A step towards quality control of herbal products. *Science, Engineering and Health Studies*, 19, Article 25050008.  
<https://doi.org/10.69598/sehs.19.25050008>
- Tiwari, R., & Shukla, A. K. (2020). Plant metabolites and their role in health benefits: A brief review. *Advance Pharmaceutical Journal*, 5(2), 47-53.  
<https://doi.org/10.31024/apj.2020.5.2.2>
- Turasan, H., Sahin, S., & Sumnu, G. (2015). Encapsulation of rosemary essential oil. *LWT-Food Science and Technology*, 64(1), 112-119. <https://doi.org/10.1016/j.lwt.2015.05.036>

- Valková, V., Ďúranová, H., Falcimaigne-Cordin, A., Rossi, C., Nadaud, F., Nesterenko, A., ... & Kačániová, M. (2022). Impact of freeze-and spray-drying microencapsulation techniques on  $\beta$ -glucan powder biological activity: A comparative study. *Foods*, 11(15), Article 2267. <https://doi.org/10.3390/foods11152267>
- Woranuch, S., & Yoksan, R. (2013). Eugenol-loaded chitosan nanoparticles: I. Thermal stability improvement of eugenol through encapsulation. *Carbohydrate Polymers*, 96(2), 578-585. <https://doi.org/10.1016/j.carbpol.2012.08.117>
- Wunnakup, T., Songsak, T., Vipunngun, N., & Sueree, L. (2024). Quantitative thin-layer chromatography analysis, antioxidant, and anti-inflammatory activities of polyherbal formulation (Ammarit-Osot) extracts. *Journal of Current Science and Technology*, 14(3), Article 65. <https://doi.org/10.59796/jcst.V14N3.2024.65>
- Yang, X. J., Glaser, R., Li, A., & Zhong, J. X. (2016a). On the aliphatic versus aromatic content of the carriers of the 'unidentified' infrared emission features. *Monthly Notices of the Royal Astronomical Society*, 462(2), 1551-1562. <https://doi.org/10.1093/mnras/stw1740>
- Yang, X., Li, A., Glaser, R., & Zhong, J. (2016b). The C-H stretching features at 3.2–3.5  $\mu\text{m}$  of polycyclic aromatic hydrocarbons with aliphatic sidegroups. *The Astrophysical Journal*, 825(1), 22. <https://doi.org/10.3847/0004-637X/825/1/22>
- Ydjedd, S., Bouriche, S., López-Nicolás, R., Sánchez-Moya, T., Frontela-Saseta, C., Ros-Berruezo, G., ... & Kati, D. E. (2017). Effect of in vitro gastrointestinal digestion on encapsulated and nonencapsulated phenolic compounds of carob (*Ceratonia siliqua* L.) pulp extracts and their antioxidant capacity. *Journal of Agricultural and Food Chemistry*, 65(4), 827-835. <https://doi.org/10.1021/acs.jafc.6b05103>
- Yoksan, R., Jirawutthiwongchai, J., & Arpo, K. (2010). Encapsulation of ascorbyl palmitate in chitosan nanoparticles by oil-in-water emulsion and ionic gelation processes. *Colloids and Surfaces B: Biointerfaces*, 76(1), 292-297. <https://doi.org/10.1016/j.colsurfb.2009.11.007>
- Yolmeh, M., & Jafari, S. M. (2017). Applications of response surface methodology in the food industry processes. *Food and Bioprocess Technology*, 10(3), 413-433. <https://doi.org/10.1007/s11947-016-1855-2>
- Zou, Y., Lu, Y., & Wei, D. (2004). Antioxidant activity of a flavonoid-rich extract of *Hypericum perforatum* L. in vitro. *Journal of Agricultural and Food Chemistry*, 52(16), 5032-5039. <https://doi.org/10.1021/jf049571r>