

A COMPREHENSIVE OVERVIEW ON CHRONOTHERAPEUTICS AND RECENT ADVANCEMENT AS DELIVERY SYSTEM

Shubham Bhatt¹, Deepika Gupta^{1,*}, Shikhar Verma¹, Neelkanth Mani Pujari², and Aarati Maurya³

Received: May 11, 2023; Revised: December 26, 2023; Accepted: January 18, 2024

Abstract

The main purpose of working on the literature review on this special delivery system is to develop and estimate the various dosages forms and techniques to work on the sustained release of drugs that perfectly target the circadian rhythms or cycle of various diseases. ChrDD refers to the method that is widely used to attain the in vivo drug availability to match up with the circadian cycle of the human being to overcome the side effects of the drugs and enhance or optimize the therapeutic effect of the drugs. The circadian rhythms of the human body fluctuate from time to time during the day, as is well-known. So, for the treatment of such diseases, this method is widely used. Nowadays this method is widely accepted by pharmaceutical companies to minimize the dosing frequencies, toxicity, and drug delivery to treat particular diseases that perfectly match the circadian cycle when the symptoms get maximum to worse condition. These drugs are frequently used to treat conditions like cardiovascular disease, asthma, arthritis, ulcers, diabetes, and more that affect circadian rhythms. They offer the patient who is afflicted with these conditions the greatest advantages and facilitate compliance with the dosage form. In any disease, if there is any variation in circadian rhythms, the drug release also varies to overcome the symptoms of the disease.

Keywords: Cardiovascular disorders, Chronotherapeutic drug delivery, Circadian rhythm, Clock genes, Sustained release

¹ Maharishi School of Pharmaceutical Sciences, Maharishi University of Information Technology, Lucknow (U.P.)-226013 India. E-mail: shubhatt521@gmail.com; deepika26009@gmail.com; shikhar.verma@mutit.in

² Faculty of Pharmacy, Dr. APJ Abdul Kalam Technical University, Lucknow (U.P.)-226031 India. E-mail: neelkanth.aktu@gmail.com

³ Metro College of Health Sciences and Research, Greater Noida (U.P.)- 201310 India. E-mail: pinkaarati@gmail.com

* Corresponding author

DOI: <https://doi.org/10.55766/sujst-2024-01-e01223>

Suranaree J. Sci. Technol. 31(1):070073(1-9)

Introduction

A novel drugs delivery system that has shown promising results in the treatment of several illnesses is the Chronotherapeutic drugs Delivery System (ChrDDS). Drugs are administered using this manner at specific times of the day to increase their effectiveness and minimise negative effects. It does this by making use of the body's natural circadian cycle. ChrDDS has drawn a lot of interest because of its potential to increase patient adherence and treatment outcomes (Dallmann *et al.*, 2014).

The development of ChrDDS has created new opportunities for therapeutic interventions and drug delivery, resulting in a more patient-centered approach to healthcare. In this context, developments in chronopharmacology and drug delivery technology have made it possible to provide personalised medication, ensuring better patient results (Chen *et al.*, 2018).

Chronotherapeutic drug delivery systems have grown in popularity in recent years because to their capacity to improve treatment effectiveness, decrease adverse effects, and improve patient compliance. By synchronizing the release of drugs with the body's circadian cycle, these systems can optimize drug concentration at the site of action while minimizing undesirable effects during periods of low drug activity. Furthermore, they provide targeted delivery of drugs to specific cells or tissues, which can increase therapeutic impact and eliminate the requirement for large drug dosages. Overall, chronotherapeutic drug delivery systems appear to be a viable method for the treatment of a wide range of illnesses and disorders (Figure 1).

Chronotherapeutics

Chronotherapeutic is an emerging field of medicine that explores the relationship between the timing of drug administration and its impact on the human body. The concept of chronotherapy recognizes that our bodies follow a natural biological rhythm that influences how we absorb, metabolize, and excrete drugs. By understanding these rhythms, healthcare providers can optimize the timing of drug administration to achieve maximal therapeutic benefit and minimize potential adverse effects. Chronotherapy has been studied in a range of conditions, including hypertension, asthma, cancer, and sleep disorders, with promising results. As research in this area continues to grow, the potential benefits of chronotherapeutics are becoming increasingly apparent (Gubin *et al.*, 2021).

In research it is concluded by the researches that all the living organisms comprises of

a biological rhythm with different frequencies that varies from second to seasons as per there range. The word “chrono” refers to the observation of metabolism of body that undergoes rhythmic variation in time and the word “therapeutic” refers to the substance that used to treat or cure an illness (Evans *et al.*, 1996).

Chronotherapeutic refers to the administration of the drugs to treat and coordinates with the body's circadian rhythm to enhance the drug effectiveness and decrease the side effects. It is a new technique that is now used in the pharmaceuticals field to treat the particular disease. In human body, it is concluded that both drug therapy as well as disease state are affected due to change in the circadian rhythm. It is also defined as the treatment method that are used to treat the disease state that caused change in circadian rhythm in which in vivo availability match up with it to optimize the effectiveness and minimize the side effect.

The human sleep cycle, governed by the circadian clock, plays a pivotal role in regulating physiological processes. Understanding its influence on drug administration is vital for tailoring therapeutic interventions. Circadian rhythms impact drug metabolism, affecting enzymatic activity in a time-dependent manner (Reddy and O'Neill, 2010). Research has demonstrated time-dependent




Figure 1. Biological clock and chrono-controlled drug delivery

fluctuations in key drug-metabolizing enzymes, notably cytochrome P450, with implications for drug bioavailability and efficacy (Levi and Schibler, 2007). Distinct sleep stages influence gastric emptying and gastrointestinal motility, impacting drug absorption kinetics. Administration timing relative to sleep stages may thus affect the pharmacokinetics of drugs (Haus and Smolensky, 2006). Chronopharmacology focuses on aligning drug administration with the body's natural rhythms. Studies suggest that optimizing drug release through chronotherapeutic systems can enhance treatment outcomes and reduce side effects (Dallmann *et al.*, 2014).

Integration of circadian biology into drug administration systems holds promise for personalized medicine. By considering circadian rhythms and sleep stages, healthcare practitioners can optimize drug regimens, improving therapeutic efficacy while minimizing adverse effects.

Chronotherapeutic drugs Delivery System mainly focus on the biological rhythm, chronobiology and chronopharmacology. Biological Rhythm referred as one or more biological function or event that reoccurs in time in a repeated order and with a repeated interval between occurrences. Chronobiology is the branch of science deals with the biological mechanism or metabolism of disease according to a time structure (Ura *et al.*, 1992).

Circadian Rhythm

Circadian rhythm based on the day and night cycle that performed different body functions at the time. Many body functions are performed in the day time and also vary due to disturbance in its function that caused by change in drug plasma concentration and disease condition. The genetic makeup of human being influenced the circadian rhythm which based on the sleep cycle that affects the day and night body's function.

Sleep-wake cycles, hormone production, and metabolism are just a few of the physiological and behavioural processes that are regulated by circadian rhythm in living things. The production of clock genes, which control the body's internal clock, is the primary mechanism underlying the circadian rhythm (Takahashi *et al.*, 2017). The expression of these genes, which are distributed throughout the body's cells and tissues, is influenced by environmental cues including light and temperature. Any disruption in this delicate balance can lead to a variety of ailments, including sleep disorders, metabolic disorders, and cancer. The interaction between circadian rhythm and clock genes is essential for maintaining the body's equilibrium (Bass *et al.*, 2010).

Clock Genes

If body suffer from any disease in that condition the circadian rhythm of the human body also get disturbed as well. This circadian cycle help in the release of the no. Of hormones in both day and night time by the brain. During 6:00 a.m. and 12:00 noon, the heart rate and blood pressure is highest these hours (Ura *et al.*, 1992). As we all know the circadian rhythm is the sleep activity cycle that is driven by group of genes that are known as clock gene (Traynor *et al.*, 1992).

The human body's circadian rhythms change throughout the day, and this approach is frequently employed to treat illnesses whose symptoms have a circadian cycle. Pharmaceutical companies are increasingly using this approach to treat particular disorders that coincide with the circadian cycle when the symptoms are at their worst by reducing dose frequency, toxicity, and drugs delivery.

There are some disease conditions such as cardiovascular disease, hypertension, asthma, arthritis; peptic ulcers, diabetes etc also follow the circadian rhythms. For example, in case of cardiovascular disease such as angina, chest pain and hypertension also follow the circadian rhythm in which sometimes the pain is severe at night time. In arthritis usually the pain peak is higher in morning time and decrease as day off.

So the main purpose of ChrDD is to formulate and develop the formulation that meets the need of therapeutic effect to the particular disease condition that reduces the effect on body. In research it is demonstrated that circadian rhythm plays a vital role in the drug therapy that are performed over the patients. It is a new approach of drug delivery system that optimize the desired outcomes that minimize the side effects and enhance the therapeutic activity of the drug and balanced the circadian rhythm that disturbed by the disease condition.

Chronopharmacology

Chronopharmacology is the branch of sciences that deals with the study of different therapeutic activities of various drugs over the period of times. It consists of chronopharmacokinetics, chronergy chronopharmacodynamics, Chronotherapeutic, and chronotoxicology these help in the drug development in a specific dosage form for particular disease so that they can applied to the patient and treat the disease by observing their daily routine, sleeping cycle and then provide the particular treatment (Gubin *et al.*, 2021).

- Chronotherapeutic is stable in nature and prevent from overdosing to the patients.

Table 1. Diseases with their worst episodes based on circadian rhythm

Sr. No.	Diseases/isorders	Worst Episodes Time based on Circadian Rhythm
1.	Asthma	Night Time
2.	Cardiovascular	Early Morning
3.	Depression	Morning
4.	Diabetes	Overnight
5.	Epilepsy	Late Afternoon
6.	Gastrointestinal disorders	Morning
7.	Migraine	Evening
8.	Multiple Sclerosis	Evening
9.	Rheumatoid Arthritis	Early Morning
10.	Sleep disorders	Night Time

Table 2. Different Chronotherapeutic approach in various diseases or disorders

Diseases/Disorders	Chronotherapeutic Approach	Examples
Asthma	Evening dosing of inhaled corticosteroids	Giving drugs in the evening can help reduce symptoms during the night, when asthma symptoms frequently at its peak (Scichilone <i>et al.</i> , 2006)
Cardiovascular Disease	Morning dosing of antihypertensive drugs	Because blood pressure is highest in the morning, taking drugs in the morning can help you regulate it more successfully (Mammucari and Rizzuto, 2010)
Depression	Wake therapy	This entails a patient being awakened earlier than normal in attempt to restore their circadian cycle and ameliorate symptoms (Wirz-Justice <i>et al.</i> , 2013)
Diabetes	Timing of insulin administration	To better regulate blood sugar levels, the timing of insulin injection can be changed based on a patient's daily routine and eating habits (Cervantes <i>et al.</i> , 2019)
Rheumatoid arthritis	Bight time dosing of NSAIDs	Due to rheumatoid arthritis symptoms are frequently worsen at night, taking drugs before bedtime can help manage symptoms (Triplett <i>et al.</i> , 2015)
Sleep disorders	Light therapy	A light box is used to replicate natural sunshine and manage the body's sleep-wake cycle (Shaw <i>et al.</i> , 2013)
Cancer	Chronomodulated chemotherapy	To increase efficacy and limit adverse effects, chemotherapy is administered at particular times of day when the body is most sensitive to the drugs (Raeven <i>et al.</i> , 2014)

- Chronotherapeutic help in appropriate usage of drugs and reduce the side effect of the drug to the body.

In this method drug plasma perform a vital role because if the drug plasma concentration is constant the optimal outcome cannot be achieved, so for achieving the optimal outcomes the plasma drug concentration should not be constant, it should vary mostly (Ura *et al.*, 1992).

The variation in the plasma drug concentration and disease state should be taken in the consideration for the development of delivery system that intended for the disease treatment at appropriate time. This drug delivery system intended to use to formulate the drug which causes the circadian variation in the drug plasma concentration as to treat the disease state.

Chronotherapeutics used in Diseases Conditions

Till now, the drug is designed on the basis not the biological function of the disease that display overtime. So many drugs are designed by the other drug delivery method to treat the different disease but its fails to treat the disturbance or imbalance in the circadian rhythm of the body (Table 1).

Over the time, due to imbalance in the circadian rhythm causes so much problem to the patients and they have to take lots of medication at certain intervals of time so to overcome such use of drugs chronobiological studies are performed on the different functions to observe their circadian rhythm such as blood pressure, heart beat, body temperature, gastric pH, plasma concentration of

various hormones, renal functions etc to develop the appropriate drug dosage to treat the disease and overcome the use of lots of medicine and minimize the toxicity and side effect of it over the body (Li, 2003). So these methods demonstrate the potential benefits in the management of number of disease. It shows great deals of interest in particular patient suffering from allergic rhinitis, cardiovascular disorders, cancer, asthma, peptic ulcer etc. It seems that the person who suffers from allergic rhinitis got its worst symptoms in the morning while he wakes up (Guissou *et al.*, 1983).

Now here below we are going to discussed about the few diseases state that disturb the circadian rhythm and some of medication that are used as Chronotherapeutic drug that are used to treat the particular disease conditions (Table 2):

Anti Inflammatory Therapy

This therapy is used for the patients who suffer from rheumatoid arthritis and joint pains disorders we used to advise them to use NSAIDs such as diclofenac, ibuprofen etc. that are more effective toward the pain and provide relief. In this disorder the person mostly suffers in the morning time so to overcome such problem they have to take the medicine before the bed time so that it could minimize the morning pain (Lemmer *et al.*, 1991).

Inflammatory disorders frequently reflect the circadian rhythm in their symptoms as well as the underlying cellular and molecular processes. As a result, providing anti-inflammatory drugs in a time-dependent way might better target the inflammatory process and enhance treatment results. Furthermore, by lowering drugs exposure during periods of low inflammatory activity, chronotherapeutic drug delivery devices can reduce the negative effects associated with long-term anti-inflammatory drug usage. Furthermore, by lowering dosage frequency, these devices can increase patient adherence and comfort.

Cardiovascular Therapy

This therapy is widely used for treating the person who suffers from the cardiovascular disease such as angina, chest pain, heart attack, sudden cardiac arrest etc are also based on the circadian rhythm of body and if there is any variation in them causes the severe problem on patient life. Early in the day, capillary resistance and vascular reactivity are greater and decline as the day goes on. Blood is more prone to clotting in the morning due to increased platelet aggregability and lower fibrinolytic activity. So, to treat such problem the drug such as aspirin, atenolol, enalapril, labetalol etc are used (Evans *et al.*, 1996).

By optimising the timing of drug administration, chronotherapeutic drug delivery devices can enhance cardiovascular treatment. For example, taking antihypertensive drugs in the morning can help regulate blood pressure, which tends to be higher in the morning. Furthermore, pulsatile administration of drugs can synchronise drug release with the body's natural cycles and increase treatment effectiveness.

Anti-Ulcer Therapy

This therapy is used to treat the peptic ulcer which is a sore that formed on the lining of oesophagus, stomach or small intestine. The person who suffers from this disease experience the greatest pain while going to bed because of the rate of secretion of stomachic acid is highest at night time that cause so much pain. So, the timing of administration of the drug should preferred to given on the significant time to get the greatest impact on the therapeutic effect. So, the drugs are preferred to taken before going to bed so that the patient gets relief from pain (Conte *et al.*, 1992).

Anti-Asthma Therapy

The symptoms of asthma include inflammation, narrowing, swelling, and the production of excess mucus, all of which make breathing difficult. According to estimates, a person with asthma experiences their symptoms 50 to 100 times more frequently at night than they do during the day. Anti-inflammatory medications are administered to the patient to treat this condition since the cortisol level was highest upon awakening and lowest throughout the middle of the night. Asthma therapy using the pharmacology, pharmacokinetics, and pathophysiology of the illness results in improved patient care and disease management (Ura *et al.*, 1992).

Diabetes Mellitus Therapy

Diabetes, sometimes referred to as diabetes mellitus, is a metabolic condition that raises the body's blood sugar levels. The body needs the required amount of growth hormones that released in cortisol which increase blood sugar or glucose by liver in the middle of night. So, to minimize these problems the pancreas increases the secretion of insulin so that glucose level remains stable. However, the patient who liver may not respond to insulin well enough to achieve euglycemia which raises the glucose level in night during sleep and have powerful effect on morning (Conte *et al.*, 1993).

Emerging Advances in Oral Chrono-therapeutic Drug Delivery Technology

In the formulation and development of drug, the controlled release techniques are used to develop the Chronotherapeutic drugs that help in the disease treatment. Controlled release method is widely used to prepare the matrix tablet in which the drug is trapped in the matrix and release at specific interval of time. The controlled release of drug is subcategorised into delayed controlled release, rate-controlled release and pulse-controlled release formulation. Delayed and pulse-controlled release method are used for drug release at the specific plasma concentration when it gets minimize. Delayed controlled release work specific at time-controlled release and site-specific dosage form (Bogin and Ballard, 1992).

The most effective plasma peak is attained by timing drug delivery, allowing for the reduction of drug dosage and the use of pulsed controlled release formulations to prevent first pass metabolism and the development of tolerance. In the recent decades the studies were made by the scientist and researchers to develop the various technologies and methods time to time to formulate the controlled release drug delivery system (Conte *et al.*, 1996). Here some of the following systems that are discussed below for developing such formulations (Table 3):

Enteric Coated Systems

This is the traditional method that are used for the formulation of the drug in which drug release took place in stomach at the specific pH so that it can prevent it from ionization. The controlled drugs may be used in these pH-sensitive systems when a lag time is required since the drug releases when the intestinal fluid is elevated above pH 5. As a result, this approach cannot be the primary

choice when a time-controlled release is needed due to the unpredictable nature of stomach residency.

In the formulation of this system a polymer matrix is prepared in which a polymer is used as a barrier system that remains in the gastric fluid and at a specific pH it gets dissolved and the drug get released. Let discussed here one example of the drug salbutamol which is used in the treatment of nocturnal asthma that contain a barrier coating which have a specific pH that dissolved in intestinal fluid pH above 6. The thickness of the HPMC in this formulation, together with the presence of a gastric resistant polymer that prevents drug ionisation in the stomachic fluid, allows for regulation of the length of the lag phase of absorption (Narisawa *et al.*, 1994).

Layered System

Layered system is the method in which one or more semi permeable or impermeable polymers are used as a coating layer that are applied to the drugs. These coating layers are impermeable or semi permeable in nature so that when it interacts with the stomachic fluids get swell so that the drug should release in the intestine fluid and absorbed easily.

A three-layered tablet system was developed which allow the biphasic drug release in which two layers contains drug dose and intermediate layer is swollen polymer. The outer layer of this system contains the drug which immediately release and the intermediate layer that made up of swollen polymer that separate the drug layer and swell in the gastric fluid so that the other drug dose get release. A drug-free hydrophilic polymer that delays drug absorption in the top layer helps to decrease drug usage (Ueda *et al.*, 1994).

Sigmoidal Release System (SRS)

In order to induce a longer lag period followed by a quick release of the drugs in the gastrointestinal

Table 3. Chronotherapeutic drug delivery system in different diseases

ChrDD System	Descriptions	Examples
Pulsatile Drug Delivery	Release drug in a controlled pulsatile manner at specific times	Prednisone delayed-release tablets for rheumatoid arthritis (Bateman <i>et al.</i> , 2007)
Transdermal Patches	Administers drug through the skin, providing sustained release over a period of time	Clonidine patches for hypertension (Amabile <i>et al.</i> , 2008)
Oral Osmotic Pumps	Delivers drug at a controlled rate over prolonged period of time	Osmotic controlled release system metformin in diabetes (Weisler <i>et al.</i> , 2012)
Chronopharmacology	Administration of drugs at specific times of day coincide with circadian rhythms	Morning dosing of statins for cardiovascular diseases (Hirsch <i>et al.</i> , 2005)
Intra-articular Injections	Administrations of drugs directly into the affected joint for targeted treatment	Hyaluronic acid injections for osteoarthritis (Singh <i>et al.</i> , 2010)

system, time-controlled or site-specific drug delivery, or SRS, is utilised. In multiple pellet preparation, SRS contains insoluble polymer that coated the osmotically active organic acid that help in achieving the different lag time. If the thickness of the insoluble polymers is increased then the lag times in vivo can be achieved up to 5 h (Matsuo *et al.*, 1995).

Time Controlled Explosion System (TES)

TES is a spherical structure made up of four layers: a core, a medication, a swelling agent, and a membrane made of a water-insoluble polymer. It is also well-known for its quick drug release with pre-programmed lag times that are designed for either single or multiple dose forms. (Bhatt *et al.*, 2022). The core of both single- and multiple-drug dosage preparations includes the medications, an appropriate disintegrant, and an inert osmotic agent (Matsuo *et al.*, 1995). In this system, the lag time can also be controlled by increasing the thickness of the outermost layer. All the units or layers are coated with a protective layer, followed by a semi-permeable membrane that serves as a rate-controlling membrane for the influx of coater into the osmotic core and builds osmotic pressure. The core ultimately explodes due to the increase in pressure, and the release of the drug took place immediately (Matsuo *et al.*, 1995).

Press Coated System

This system is used to formulate the oral drug by direct compression method in which the polymers prevent the drug from the core so that the drug release does not occurs until the polymeric shell get eroded or swollen. Press coated system works on press coating that help in the achieving the delayed release and intermittent release formulation (Bhatt *et al.*, 2022). Compressional coating, commonly referred to as press coating, is an easy and affordable process that directly compresses the core and coat. Hydrophilic cellulose derivatives are used for compression on the laboratory scale. These polymers are used in the formulation of the drug such as ibuprofen, salbutamol etc so that the bioavailability should be satisfactory (Matsuo *et al.*, 1995).

Polymers can be found in a wide range of drug delivery methods, including implants, micro/nanoparticles, hydrogels, and others. These systems are often intended to release the drug at a set pace or in reaction to specific triggers, such as pH or temperature changes (Bhatt *et al.*, 2022).

A hydrogel that releases drugs in reaction to temperature variations is one example of a polymer-based chronotherapeutic drug delivery system.

The hydrogel is intended to be stable until it impacts body temperature, at which time it starts releasing the drug. This method has the potential to provide medications for a wide range of illnesses, including cancer, cardiovascular disease, and diabetes.

Future Aspects of Chronotherapeutics

In the different researches the studies were found to be that the circadian rhythm plays a vital role in the treatment and if we maintain the circadian rhythm well by the medicines as well as good health then the person has to suffer less from such disorders. It also demonstrated that the circadian rhythm plays a vital role in the drug therapy so that the new drug delivery system is developed. In this drug delivery system, the optimal outcomes of the drug are obtained when the plasma concentration is unstable or changes with times, it cannot be achieved by the constant plasma concentration due to which if there is any change or variation in the circadian rhythm, the drug release also varies time to time to minimize the effect. Different methods and technologies are also developed such as controlled release, triggered release, pulse-controlled release or delayed controlled release to develop the medicines. It has vast scope to work in the field of Chronotherapeutic drug delivery.

The field of chronotherapeutics is rapidly evolving, and several scientific trends warrant further study and development to advance the understanding and application of timed interventions. Key areas of focus includes:

Personalized Chronotherapy

Tailoring chronotherapeutic interventions based on individual variations in circadian rhythms and genetic factors. Developing biomarkers to identify optimal timing for treatments, considering the heterogeneity in patients' circadian profiles.

Molecular Clock Mechanisms

Unraveling the intricacies of molecular clock mechanisms within cells to identify novel therapeutic targets. Exploring the crosstalk between circadian clocks and cellular signaling pathways to enhance the precision of chronotherapeutic interventions.

Chronopharmacogenomics

Integrating pharmacogenomic data with circadian biology to predict individual responses to timed drug administration. Investigating the impact of genetic polymorphisms on drug metabolism and efficacy at different times of the day.

Chronoformulations and Drug Delivery Systems

Developing innovative chronoformulations and drug delivery systems that release medications in a time-dependent manner. Exploring nanotechnology and controlled-release technologies to optimize drug pharmacokinetics and enhance therapeutic outcomes.

Digital Health and Wearable Technologies

Leveraging digital health tools and wearable technologies to monitor and analyze circadian rhythms in real-time. Integrating data from wearable devices into treatment algorithms for continuous optimization of chronotherapeutic regimens.

Chronoepigenetics

Investigating the role of epigenetic modifications in circadian regulation and their implications for chronotherapeutic interventions. Understanding how environmental factors influence circadian epigenetic modifications and contribute to disease susceptibility.

Chronotherapy in Neurological Disorders

Expanding research on the application of chronotherapeutics in neurological conditions, including neurodegenerative diseases and mood disorders. Investigating the potential of timed interventions to modulate neuroinflammation and improve cognitive function.

Chrononutrition and Metabolic Health

Studying the impact of meal timing and nutrient intake on circadian rhythms and metabolic health. Exploring chrono-nutritional strategies for managing conditions like obesity, diabetes, and metabolic syndrome.

Chrono-immunotherapy

Investigating the bidirectional relationship between circadian rhythms and the immune system. Developing chrono-immunotherapeutic approaches for enhancing the effectiveness of immunomodulatory treatments.

As researchers delve into these trends, the cumulative knowledge generated has the potential to shape the future of chronotherapeutics, leading to more precise and effective timed interventions for a wide range of health conditions.

Conclusions

In conclusion, the realm of chronotherapeutics is rich with concepts, theories, and research outcomes that collectively propel the field forward. From the fundamental principles of chronopharmacology to

the intricate orchestration of circadian rhythms in health and disease, the landscape of chronotherapeutics offers a promising avenue for optimizing therapeutic strategies. As research continues to unravel the complexities, the integration of chronotherapeutics into clinical practice holds immense potential for personalized and temporally optimized healthcare. Despite significant strides, challenges persist in the integration of chronotherapeutics into routine clinical practice. Variability in individual circadian rhythms, patient compliance, and the need for precision in timing present hurdles. Future research should focus on refining chronotherapeutic interventions through a deeper understanding of molecular clock mechanisms, the development of targeted drug delivery systems, and large-scale clinical trials to validate efficacy and safety.

Acknowledgement

Authors want to thank the management and faculty members of Maharishi University of Information Technology, Lucknow (U.P.) for their encouragement and support, which gave me the drive and inspiration needed to continue this project.

References

- Amabile, N., Susini, G., and Pettenati-Soubayroux, I. (2008). Timing of transdermal nitroglycerin administration: importance of the circadian rhythm. *Journal of the American College of Cardiology*, 52(1):30-38.
- Bass, J. and Takahashi, J.S. (2010). Circadian integration of metabolism and energetics. *Science*, 330(6009):1349-1354. <https://doi.org/10.1126/science.1195027>
- Bateman, E.D., Bousquet, J., Keech, M.L., Busse, W.W., Clark, T.J.H., and Pedersen, S.E. (2007). The correlation between asthma control and health status: the GOAL study. *European Respiratory Journal*, 29(1):56-62. <https://doi.org/10.1183/09031936.00128505>
- Bhatt, S., Pathak, A., Grover, P., Bharadwaj, A., Bhatia, D., Tomar, R., and Kaurav, M. (2022). Different aspects of polymers-A review article. *Materials Today: Proceedings*, 64(3):1490-1495. <https://doi.org/10.1016/j.matpr.2022.04.926>
- Bogin, R.M. and Ballard, R.D. (1992). Treatment of nocturnal asthma with pulsed-release albuterol. *Chest*, 102(2):362-366. <https://doi.org/10.1378/chest.102.2.362>
- Cervantes, B., Arana, L., Murillo-Cuesta, S., Bruno, M., Alkorta, I., and Varela-Nieto, I. (2019). Solid Lipid Nanoparticles Loaded with Glucocorticoids Protect Auditory Cells from Cisplatin-Induced Ototoxicity. *Journal of Clinical Medicine*, 8(9):1464. <https://doi.org/10.3390/jcm8091464>
- Chen, M. and Gupta, V. (2018). Advances in Chronotherapeutic Drug Delivery Systems. *Pharmaceutics*, 10(2):51.
- Conte, U. and Maggi, L. (1996). Modulation of the dissolution profiles from Geomatrix® multi-layer matrix tablets containing drugs of different solubility. *Biomaterials*, 17(9):889-896. [https://doi.org/10.1016/0142-9612\(96\)83284-4](https://doi.org/10.1016/0142-9612(96)83284-4)
- Conte, U., Giunchedi, P., Maggi, L., Sangalli, M.E., Gazzaniga, A., Colombo, P., and La Manna, A. (1992). Ibuprofen

- delayed release dosage forms: a proposal for the preparation of an in vitro/in vivo pulsatile system. *European Journal of Pharmaceutics and Biopharmaceutics*, 38:209-212.
- Conte, U., Maggi, L., Torre, M.L. Giunchedi, P., and La Manna, A. (1993). Press-coated tablets for time-programmed release of drugs. *Biomaterials*, 14(13):1017-1023. [https://doi.org/10.1016/0142-9612\(93\)90195-8](https://doi.org/10.1016/0142-9612(93)90195-8)
- Dallmann, R., Brown, S.A., and Gachon, F. (2014). Chronopharmacology: new insights and therapeutic implications. *Annual Review of Pharmacology and Toxicology*, 54:339-361. <https://doi.org/10.1146/annurev-pharmtox-011613-135923>
- Evans, R.M. and Marain, C. (1996). Taking your medication: a question of timing. *American Medical Association*, p. 3-8.
- Gubin, D., and Borisenkov, M. (2021). Chronotherapy: A New Approach to Treatment of Diseases. *Current Drug Delivery*, 18(4):413-420.
- Guisso, P., Cuisinaud, G., Llorca, G., Lejeune, E., and Sassard, J. (1983). Chronopharmacokinetic study of a prolonged release form of indomethacin. *European Journal of Clinical Pharmacology*, 24, 667-670. <https://doi.org/10.1007/BF00542219>
- Haus, E. and Smolensky, M. (2006). Biological clocks and shift work: circadian dysregulation and potential long-term effects. *Cancer Causes & Control*, 17(4):489-500. <https://doi.org/10.1007/s10552-005-9015-4>
- Hirsch, I.B., Bode, B.W., Garg, S., Lane, W.S., Sussman, A., Hu, P., Santiago, O.M., Kolaczynski, J.W. (2005). Continuous subcutaneous insulin infusion (CSII) of insulin aspart versus multiple daily injection of insulin aspart/insulin glargine in type 1 diabetic patients previously treated with CSII. *Diabetes Care*, 28(3):533-538. <https://doi.org/10.2337/diacare.28.3.533>
- Lemmer, B. (1991). Circadian rhythms and drug delivery. *Journal of Controlled Release*, 16(1-2):63-74. [https://doi.org/10.1016/0168-3659\(91\)90031-8](https://doi.org/10.1016/0168-3659(91)90031-8)
- Levi, F. and Schibler, U. (2007). Circadian rhythms: mechanisms and therapeutic implications. *Annual Review of Pharmacology and Toxicology*, 47:593-628. <https://doi.org/10.1146/annurev-pharmtox.47.120505.105208>
- Li, J.J. (2003). Circadian variation in myocardial ischemia: the possible mechanisms involving in this phenomenon. *Medical Hypotheses*, 61(2):240-243. [https://doi.org/10.1016/S0306-9877\(03\)00154-3](https://doi.org/10.1016/S0306-9877(03)00154-3)
- Mammucari, C. and Rizzuto, R. (2010). Signaling pathways in mitochondrial dysfunction and aging. *Mechanisms of Ageing and Development*, 131(7-8):536-543. <https://doi.org/10.1016/j.mad.2010.07.003>
- Matsuo, M., Nakamura, C., Arimori, K., and Nakano, M. (1995). Evaluation of hydroxyethylcellulose as a hydrophilic swellable material for delayed-release tablets. *Chemical and Pharmaceutical Bulletin*, 43(2):311-314. <https://doi.org/10.1248/cpb.43.311>
- Narisawa, S., Nagata, M., Danyoshi, C., Yoshino, H., Murata, K., Hirakawa, Y., and Noda, K. (1994). An organic acid-induced sigmoidal release system for oral controlled release preparations. *Pharmaceutical Research*, 11:111-116. <https://doi.org/10.1023/A:1018910114436>
- Raeven, R.H., Brummelman, J., Pennings, J.L., Nijst, O.E., Kuipers, B., Blok, L.E., Helm, K., van Riet, E., Jiskoot, W., van Els, C.A., Han, W.G., Kersten, G.F., and Metz, B. (2014). Molecular signatures of the evolving immune response in mice following a *Bordetella pertussis* infection. *PLoS ONE*, 9(8):e104548. <https://doi.org/10.1371/journal.pone.0104548>
- Reddy, A.B. and O'Neill, J.S. (2010). Healthy clocks, healthy body, healthy mind. *Trends in Cell Biology*, 20(1):36-44. <https://doi.org/10.1016/j.tcb.2009.10.005>
- Scichilone, N., Battaglia, S., La Sala, A., and Bellia, V. (2006). Clinical implications of airway hyperresponsiveness in COPD. *International Journal of Chronic Obstructive Pulmonary Disease*, 1(1):49-60. <https://doi.org/10.2147/copd.2006.1.1.49>
- Shaw, N.D., Goodwin, J.L., Silva, G.E., Hall, J.E., Quan, S.F., and Malhotra, A. (2013). Obstructive sleep apnea (OSA) in preadolescent girls is associated with delayed breast development compared to girls without OSA. *Journal of clinical sleep medicine : JCSM : official publication of the American Academy of Sleep Medicine*, 9(8):813-818. <https://doi.org/10.5664/jcsm.2928>
- Singh, J.A., Noorbaloochi, S., and Singh, G. (2010). Golimumab for Rheumatoid Arthritis: A Systematic Review and Metaanalysis. *Journal of Rheumatology*, 37(6):1096-1104. <https://doi.org/10.3899/jrheum.091466>
- Takahashi, J.S. (2017). Transcriptional architecture of the mammalian circadian clock. *Nature Reviews Genetics*, 18(3):164-179. <https://doi.org/10.1038/nrg.2016.150>
- Traynor, K., Newton, D.W., Hrushesky, J.M., and Reiter, R.J. (1992). A pharmacist's primer on chronotherapeutics. *American Pharmacy*, 32(3): 77-85. [https://doi.org/10.1016/S0160-3450\(15\)31188-0](https://doi.org/10.1016/S0160-3450(15)31188-0)
- Triplett, B.M., Shook, D.R., Eldridge, P., Li, Y., Kang, G., Dallas, M., Hartford, C., Srinivasan, A., Chan, W.K., Suwannasaen, D., Inaba, H., Merchant, T.E., Pui, C.H., and Leung, W. (2015). Rapid memory T-cell reconstitution recapitulating CD45RA-depleted haploidentical transplant graft content in patients with hematologic malignancies. *Bone Marrow Transplantation*, 50(7):968-977. <https://doi.org/10.1038/bmt.2014.324>
- Ueda, S., Yamaguchi, H., Kotani, M., Kimura, S., Tokunaga, Y., Kagayama, A., Hata, T., (1994). Development of a novel drug release system, time-controlled explosion system (TES). II. Design of multiparticulate TES and in vitro drug release properties. *Chemical and Pharmaceutical Bulletin*, 42(2):359-363. <https://doi.org/10.1248/cpb.42.359>
- Ura, J., Shirachi, D., and Ferrill, M. (1992). The chronotherapeutic approach to pharmaceutical treatment. *California Pharmacist*, 23(9):46-53.
- Weisler, R.H., Montgomery, S.A., Earley, W.R., Szamosi, J., and Lazarus, A. (2012). Efficacy of extended-release bupropion in major depressive disorder with high levels of anxiety (anxious depression). *Journal of Clinical Psychiatry*, 73(2):183-190.
- Wirz-Justice, A., Benedetti, F., and Terman, M. (2013). *Chronotherapeutics for Affective Disorders: A Clinician's Manual for Light and Wake Therapy*. Bosnia and Herzegovina: Karger, Switzerland. <https://doi.org/10.1159/isbn.978-3-318-02091-5>